# Summary of Anticoagulant Monitoring Guidelines

(Pharmacists may order labs more often if needed.)

#### warfarin (Coumadin)

#### INR:

Initiation Dosing Protocol - baseline unless within past 24 hours and then daily.
Chronic Dosing Protocol - baseline unless within past 24 hours and then at least weekly.
Monitoring Protocol - baseline unless within past 24 hours and then daily or per MD or Pharmacist order.

#### <u>CBC:</u>

Initiation Dosing Protocol - baseline unless within past 24 hours and then at least weekly. Chronic Dosing Protocol - baseline unless within past 24 hours and then at least weekly. Monitoring Protocol - baseline unless within past 24 hours and then at least weekly.

## enoxaparin (Lovenox)

#### <u>CBC:</u>

Baseline unless within past 24 hours and then every 3 days. If platelet count and hgb are stable x 2 weeks and there are no signs of bleeding, then weekly.

<u>SCr:</u>

Baseline unless within past 24 hours and then every 3 days. If stable x 1 week, then weekly.

# fondaparinux (Arixtra)

<u>CBC:</u>

Baseline unless within past 24 hours and then weekly.

<u>SCr:</u>

Baseline unless within past 24 hours and then weekly.

## **Prophylactic SQ heparin**

<u>CBC:</u>

Baseline unless within past 24 hours and then every 3 days. If platelet count and hgb are stable x 2 weeks and there are no signs of bleeding, then weekly.

# Oral anticoagulants - apixaban (Eliquis), dabigatran (Pradaxa), and rivaroxaban (Xarelto)

<u>CBC:</u>

Baseline unless within past 24 hours and then weekly.

<u>SCr:</u>

Baseline unless within past 24 hours and then weekly.