Entering PD Exchange Bags via PowerChart

Search for PD Exchange. Options below. Select appropriate administration method and strength.

PD Exchange Cycler 1.5% PD Exchange Cycler 1.5% 8000 mL, Misc, Intraperitoneal, Once PD Exchange Cycler 2.5% PD Exchange Cycler 2.5% 8000 mL, Misc, Intraperitoneal, Once PD Exchange Cycler 4.25% PD Exchange Cycler 4.25% 8000 mL, Misc, Intraperitoneal, Once PD Exchange Manifold 1.5% PD Exchange Manifold 1.5% 2000 mL, Misc, Intraperitoneal, QID, dose(s)/time(s) PD Exchange Manifold 2.5% PD Exchange Manifold 2.5% 2000 mL, Misc, Intraperitoneal, QID, dose(s)/time(s) PD Exchange Manifold 4.25% PD Exchange Manifold 4.25% 2000 mL, Misc, Intraperitoneal, QID, dose(s)/time(s) PD Exchange Manual 1.5% PD Exchange Manual 1.5% 2000 mL, Misc, Intraperitoneal, QID, dose(s)/time(s) PD Exchange Manual 2.5% PD Exchange Manual 2.5% 2000 mL, Misc, Intraperitoneal, QID, dose(s)/time(s) PD Exchange Manual 4.25% PD Exchange Manual 4.25% 2000 mL, Misc, Intraperitoneal, QID, dose(s)/time(s)

Select Intermittent and OK

P Infusion Type X

How would you like to give PD Exchange Manual 2.5%?

Give As:		
O Continuous infusion		
OIntermittent infusion		
	OK	Cancel

Add Order Comments as indicated by provider

⊿ Medications						
	Peritoneal Dialysis	; Order	6/22/2022 11:00 E	DT 8,000 mL, Misc	, Intraperitoneal, Or	nce, Routine, Start date: 6/2
	(Dianeal Low Calc	ium		AmbuFlex III vi	a Cycler. Same as D	ianeal PD-2 with 1.5% Dex
Details for Peritone	al Dialysis (E	Dianeal Lov	w Calcium)	-D1.5% (PL) Exchange	e Cycler 1.5%)
Petails := 📴 Ingredie	ent Details 🕞 Diag	noses				
Ingredients	Dose	Rate	Infuse Over	Frequency	Duration	
× PD Exchange Cycler 1.5%	8,000 mL 🗧 🗧	6		Once 🗸		
1						
	Total Volume					
	8,000 mL					
Order comments						
AmbuFlex III via Cycler, Sa	me as Dianeal PD-2 w	ith 1.5% Dextrose.				
PD 1.5%, 2000 mL, cycler, 4	4 exchanges over 9 ho	urs. Add KCI 2 mE	q per liter of dialysa	ate.		
1						



Add # doses. (For SNF, may need to remove the "dose(s)/time(s).)

On Details tab, review / edit the QID frequency or administration times. Use the down arrow to change the frequency. Click on the field, then click on the three dots to edit the administration times.



[™] Reschedule



Dextrose 2.5% with Electrolytes (Dianeal Low Calcium) intraperitoneal solution (PD Exchange Manual 2.5%)

PHARMACY INFO

Pharmacy will place appropriate order or orders needed based on the Order Comments.

Tidbits on Order Entry for Cycler in PharmNet

- Infuse Over: 1 Hour
- If no Additives, only one order is needed
- If Additives:
 - One order per bag prepared is needed. This is so each bag can be labeled appropriately with the volume of bag and any additive amounts.
 - Volume of PD bag being used needs to be the Volume for the order ie, if 6 L bag being used, the PD volume would be 6 L. This is true even if the order is for 8000 mL
 - Additive amount needs to be adjusted based on the size of the bag ie, KCl 2 mEq per liter would be 12 mEq per 6 L bag