

The Single Shared System for Clozapine No Blood, No DrugTM

Recommended Monitoring Frequency and Clinical Decisions by ANC Level

ANC Level	Treatment Recommendation	ANC Monitoring
Normal Range for a New Patient • General Population (ANC ≥ 1500/µL) BEN POPULATION	 Initiate treatment If treatment interrupted: < 30 days, continue monitoring as before ≥ 30 days, monitor as if new patient 	 Weekly from initiation to 6 months Every 2 weeks from 6 to 12 months Monthly after 12 months
 BEN Population (ANC ≥ 1,000/μL) Obtain at least two baseline ANC levels before initiating treatment 	Discontinuation for reasons other than neutropenia	See Section 2.4 of the full Prescribing Information
Mild Neutropenia (1000 to 1499/μL)*	• Continue treatment	GENERAL POPULATION • Three times weekly until ANC ≥ 1500/μL • Once ANC ≥ 1500/μL, return to patient's last "Normal Range" ANC monitoring interval**
	 BEN POPULATION Mild Neutropenia is normal range for BEN population, continue treatment Obtain at least two baseline ANC levels before initiating treatment If treatment interrupted < 30 days, continue monitoring as before ≥ 30 days, monitor as if new patient 	 BEN POPULATION Weekly from initiation to 6 months Every 2 weeks from 6 to 12 months Monthly after 12 months
	Discontinuation for reasons other than neutropenia	See Section 2.4 of the full Prescribing Information
Moderate Neutropenia (500 to 999/µL)*	 GENERAL POPULATION Recommend hematology consultation Interrupt treatment for suspected clozapine induced neutropenia Resume treatment once ANC normalizes to ≥ 1000/μL 	 GENERAL POPULATION Daily until ANC ≥ 1000/μL, then Three times weekly until ANC ≥ 1500/μL Once ANC ≥ 1500/μL, check ANC weekly for 4 weeks, then return to patient's last "Normal Range" ANC monitoring interval**
	BEN POPULATION Recommend hematology consultation Continue treatment	 BEN POPULATION Three times weekly until ANC ≥ 1000/μL or ≥ patient's known baseline. Once ANC ≥ 1000/μL or patient's known baseline, then check ANC weekly for 4 weeks, then return to patient's last "Normal BEN Range" ANC monitoring interval**
Severe Neutropenia (less than 500/µL)*	GENERAL POPULATION Recommend hematology consultation Interrupt treatment for suspected clozapine induced neutropenia Do not rechallenge unless prescriber determines benefits outweigh risks	 GENERAL POPULATION Daily until ANC ≥ 1000/μL Three times weekly until ANC ≥ 1500/μL If patient rechallenged, resume treatment as a new patient under "Normal Range" monitoring once ANC ≥ 1500/μL
	BEN POPULATION Recommend hematology consultation Interrupt treatment for suspected clozapine induced neutropenia Do not rechallenge unless prescriber determines benefits outweigh risks	 BEN POPULATION Daily until ANC ≥ 500/μL Three times weekly until ANC ≥ patients established baseline If patient rechallenged, resume treatment as a new patient under "Normal Range" monitoring once ANC ≥1000/μL or at patient's baseline

^{*} Confirm all initial reports of ANC less than 1500/µL (ANC < 1000/µL for BEN patients) with a repeat ANC measurement within 24 hours

^{**} If clinically appropriate