

Emergency Department and Hospital Management of Pit Viper Snakebite (Rattlesnakes, copperheads and cottonmouths)

!!!Antivenom must be ordered by an attending physician!!!
!!!Admit patient to the critical care medicine service post antivenom administration!!!

IV Analgesics

- Morphine 4mg IV every 4 hours PRN severe pain [ERX: 175930]
- Hydromorphone 0.2 to 1 mg IV every 2 to 3 hours PRN severe pain [ERX: 175878]
- Fentanyl 50mcg IV q2h PRN severe pain [ERX: 3037]

Fluids

- Sodium Chloride 0.9% Bolus 1000mL IV once [ERX: 400291]
- Lactated Ringers 1000mL IV once [EXR: 400296]

Tetanus

- Diph, pertuss(ace), tet vac(PF) (Adacel) 0.5mL IM once [ERX: 520179]

Labs

- Fibrinogen
- Protimes-INR
- CBC
- D-dimer
- PTT

Antivenom

- For adult patients with confirmed copperheads, cottonmouth/water moccasin

Initial dose: 4 vials. Consider 8 vials for patients in shock or serious active bleeding. May repeat additional 4 vials if control is not achieved with initial dose. Max initial dose: 12 vials

- Initial Dose
 - Crotalidae polyvalent immune fab (CROFAB) 4 vials in 0.9% sodium chloride 250 mL IVPB [ERX: 520136]
- Maintenance Dose
 - Crotalidae polyvalent immune fab (CROFAB) 2 vials in sodium chloride 0.9% 250 mL IVPB [500325]
 - Ordering instruction: 2 vials every 6 hours for 3 doses
- For adult patients with rattlesnake or unknown snake envenomation
 - Initial Dose
 - Antivenin Crotalidae (equine) (ANAVIP) 10 vials in 0.9% sodium chloride 250mL IVPB [ERX: 501030]
 - Maintenance Dose
 - Antivenin Crotalidae (equine) (ANAVIP) 4 vials in 0.9% sodium chloride 250mL IVPB [ERX: 501031]